

## HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use AUSTEDO safely and effectively. See full prescribing information for AUSTEDO.

**AUSTEDO™ (deutetrabenazine) tablets, for oral use**  
Initial U.S. Approval: 2017

### WARNING: DEPRESSION AND SUICIDALITY

See full prescribing information for complete boxed warning.

- Increases the risk of depression and suicidal thoughts and behavior (suicidality) in patients with Huntington's disease (5.2)
- Balance risks of depression and suicidality with the clinical need for treatment of chorea when considering the use of AUSTEDO (5.2)
- Monitor patients for the emergence or worsening of depression, suicidality, or unusual changes in behavior (5.2)
- Inform patients, caregivers and families of the risk of depression and suicidality and instruct to report behaviors of concern promptly to the treating physician (5.2)
- Exercise caution when treating patients with a history of depression or prior suicide attempts or ideation (5.2)
- AUSTEDO is contraindicated in patients who are suicidal, and in patients with untreated or inadequately treated depression (4, 5.2)

### INDICATIONS AND USAGE

AUSTEDO is a vesicular monoamine transporter 2 (VMAT2) inhibitor indicated for the treatment of chorea associated with Huntington's disease (1)

### DOSAGE AND ADMINISTRATION

- The starting dose is 6 mg once daily. Titrate up at weekly intervals by 6 mg per day to a tolerated dose that reduces chorea, up to a maximum recommended daily dosage of 48 mg (24 mg twice daily) (2.1)
- Administer total daily dosages of 12 mg or above in two divided doses (2.1)
- Administer with food (2.1)
- Swallow tablets whole; do not chew, crush, or break (2.1)
- If switching patients from tetrabenazine, discontinue tetrabenazine and initiate AUSTEDO the following day. See full prescribing information for recommended conversion table (2.2)
- Maximum recommended dosage of AUSTEDO in poor CYP2D6 metabolizers is 36 mg per day (i.e., 18 mg twice daily) (2.4, 8.7)

## FULL PRESCRIBING INFORMATION: CONTENTS\*

### WARNING: DEPRESSION AND SUICIDALITY

#### 1 INDICATIONS AND USAGE

#### 2 DOSAGE AND ADMINISTRATION

- 2.1 Dosing Information
- 2.2 Switching Patients from Tetrabenazine (XENAZINE®) to AUSTEDO
- 2.3 Dosage Adjustment with Strong CYP2D6 Inhibitors
- 2.4 Dosage Adjustment in Poor CYP2D6 Metabolizers
- 2.5 Discontinuation and Interruption of Treatment

#### 3 DOSAGE FORMS AND STRENGTHS

#### 4 CONTRAINDICATIONS

#### 5 WARNINGS AND PRECAUTIONS

- 5.1 Clinical Worsening and Adverse Events
- 5.2 Depression and Suicidality
- 5.3 Neuroleptic Malignant Syndrome (NMS)
- 5.4 Akathisia, Agitation, and Restlessness
- 5.5 Parkinsonism
- 5.6 Sedation and Somnolence
- 5.7 QTc Prolongation
- 5.8 Hyperprolactinemia
- 5.9 Binding to Melanin-Containing Tissues

#### 6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience

#### 7 DRUG INTERACTIONS

- 7.1 Strong CYP2D6 Inhibitors
- 7.2 Reserpine
- 7.3 Monoamine Oxidase Inhibitors (MAOIs)

## FULL PRESCRIBING INFORMATION

### WARNING: DEPRESSION AND SUICIDALITY

AUSTEDO can increase the risk of depression and suicidal thoughts and behavior (suicidality) in patients with Huntington's disease. Anyone considering the use of AUSTEDO must balance the risks of depression and suicidality with the clinical need for treatment of chorea. Closely monitor patients for the emergence or worsening of depression, suicidality, or unusual changes in behavior. Patients, their caregivers, and families should be informed of the risk of depression and suicidality and should be instructed to report behaviors of concern promptly to the treating physician.

Particular caution should be exercised in treating patients with a history of depression or prior suicide attempts or ideation, which are increased in frequency in Huntington's disease. AUSTEDO is contraindicated in patients who are suicidal, and in patients with untreated or inadequately treated depression [see *Contraindications (4) and Warnings and Precautions (5.2)*].

## DOSAGE FORMS AND STRENGTHS

Tablets: 6 mg, 9 mg, and 12 mg (3)

## CONTRAINDICATIONS

- Suicidal, or untreated/inadequately treated depression (4, 5.2)
- Hepatic impairment (4, 8.6, 12.3)
- Taking MAOIs, reserpine, or tetrabenazine (XENAZINE®) (4, 7.2, 7.3, 7.7)

## WARNINGS AND PRECAUTIONS

- Neuroleptic Malignant Syndrome (NMS): Discontinue if this occurs (5.3, 7.4)
- Akathisia, agitation, restlessness, and parkinsonism: Reduce dose or discontinue if this occurs (5.4, 5.5)
- Sedation/somnolence: May impair the patient's ability to drive or operate complex machinery (5.6)

## ADVERSE REACTIONS

Most common adverse reactions (>8% of AUSTEDO-treated patients and greater than placebo) were: somnolence, diarrhea, dry mouth, and fatigue (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Teva Pharmaceuticals at 1-888-483-8279 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

## DRUG INTERACTIONS

- Concomitant use of strong CYP2D6 inhibitors: Maximum recommended dose of AUSTEDO is 36 mg per day (18 mg twice daily) (2.3, 7.1)
- Alcohol or other sedating drugs: May have additive sedation and somnolence (7.5)

## USE IN SPECIFIC POPULATIONS

Pregnancy: Based on animal data, may cause fetal harm (8.1)

See 17 for PATIENT COUNSELING INFORMATION and Medication Guide.

Revised: 4/2017

- 7.4 Neuroleptic Drugs
- 7.5 Alcohol or Other Sedating Drugs
- 7.6 Drugs that Cause QTc Prolongation
- 7.7 Tetrabenazine

#### 8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Hepatic Impairment
- 8.7 Poor CYP2D6 Metabolizers

#### 10 OVERDOSAGE

#### 11 DESCRIPTION

#### 12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

#### 13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

#### 14 CLINICAL STUDIES

#### 16 HOW SUPPLIED/STORAGE AND HANDLING

- 16.1 How Supplied
- 16.2 Storage

#### 17 PATIENT COUNSELING INFORMATION

\*Sections or subsections omitted from the full prescribing information are not listed.

## 1 INDICATIONS AND USAGE

AUSTEDO™ is indicated for the treatment of chorea associated with Huntington's disease.

## 2 DOSAGE AND ADMINISTRATION

### 2.1 Dosing Information

The dose of AUSTEDO is determined individually for each patient based on reduction of chorea and tolerability. When first prescribed to patients who are not being switched from tetrabenazine (a related VMAT2 inhibitor), the recommended starting dose of AUSTEDO is 6 mg administered orally once daily.

- The dose of AUSTEDO may be increased at weekly intervals in increments of 6 mg per day to a maximum recommended daily dosage of 48 mg.
- Administer total daily dosages of 12 mg or above in two divided doses.
- Administer AUSTEDO with food [see *Clinical Pharmacology (12.3)*].
- Swallow AUSTEDO whole. Do not chew, crush, or break tablets.

**2.2 Switching Patients from Tetrabenazine (XENAZINE®) to AUSTEDO**

Discontinue tetrabenazine (XENAZINE®) and initiate AUSTEDO the following day. The recommended initial dosing regimen of AUSTEDO in patients switching from tetrabenazine (XENAZINE®) to AUSTEDO is shown in Table 1.

**Table 1. Recommended Initial Dosing Regimen when Switching from Tetrabenazine (XENAZINE®) to AUSTEDO**

Current tetrabenazine daily dosage	Initial regimen of AUSTEDO
12.5 mg	6 mg once daily
25 mg	6 mg twice daily
37.5 mg	9 mg twice daily
50 mg	12 mg twice daily
62.5 mg	15 mg twice daily
75 mg	18 mg twice daily
87.5 mg	21 mg twice daily
100 mg	24 mg twice daily

After patients are switched to AUSTEDO, the dose may be adjusted at weekly intervals [see *Dosage and Administration* (2.1)].

**2.3 Dosage Adjustment with Strong CYP2D6 Inhibitors**

In patients receiving strong CYP2D6 inhibitors (e.g., quinidine, antidepressants such as paroxetine, fluoxetine, and bupropion), the total daily dosage of AUSTEDO should not exceed 36 mg (maximum single dose of 18 mg) [see *Drug Interactions* (7.1) and *Clinical Pharmacology* (12.3)].

**2.4 Dosage Adjustment in Poor CYP2D6 Metabolizers**

In patients who are poor CYP2D6 metabolizers, the total daily dosage of AUSTEDO should not exceed 36 mg (maximum single dose of 18 mg) [see *Use in Specific Populations* (8.7)].

**2.5 Discontinuation and Interruption of Treatment**

Treatment with AUSTEDO can be discontinued without tapering. Following treatment interruption of greater than one week, AUSTEDO therapy should be re-titrated when resumed. For treatment interruption of less than one week, treatment can be resumed at the previous maintenance dose without titration.

**3 DOSAGE FORMS AND STRENGTHS**

AUSTEDO tablets are available in the following strengths:

- The 6 mg tablets are round, purple-coated tablets, with “SD” over “6” printed in black ink on one side.
- The 9 mg tablets are round, blue-coated tablets, with “SD” over “9” printed in black ink on one side.
- The 12 mg tablets are round, beige-coated tablets, with “SD” over “12” printed in black ink on one side.

**4 CONTRAINDICATIONS**

AUSTEDO is contraindicated in patients:

- Who are suicidal, or in patients with untreated or inadequately treated depression [see *Warnings and Precautions* (5.2)].
- With hepatic impairment [see *Use in Specific Populations* (8.6), *Clinical Pharmacology* (12.3)].
- Taking monoamine oxidase inhibitors (MAOIs). AUSTEDO should not be used in combination with an MAOI, or within 14 days of discontinuing therapy with an MAOI [see *Drug Interactions* (7.3)].
- Taking reserpine. At least 20 days should elapse after stopping reserpine before starting AUSTEDO [see *Drug Interactions* (7.2)].
- Taking tetrabenazine (XENAZINE®) [see *Drug Interactions* (7.7)].

**5 WARNINGS AND PRECAUTIONS****5.1 Clinical Worsening and Adverse Events**

Huntington's disease is a progressive disorder characterized by changes in mood, cognition, chorea, rigidity, and functional capacity over time. VMAT2 inhibitors, including AUSTEDO, may cause a worsening in mood, cognition, rigidity, and functional capacity.

Prescribers should periodically re-evaluate the need for AUSTEDO in their patients by assessing the effect on chorea and possible adverse effects, including sedation/somnolence, depression and suicidality, parkinsonism, akathisia, restlessness, and cognitive decline. It may be difficult to distinguish between adverse reactions and progression of the underlying disease; decreasing the dose or stopping the drug may help the clinician to distinguish between the two possibilities. In some patients, the underlying chorea itself may improve over time, decreasing the need for AUSTEDO.

**5.2 Depression and Suicidality**

Patients with Huntington's disease are at increased risk for depression, and suicidal ideation or behaviors (suicidality). AUSTEDO may increase the risk for suicidality in patients with Huntington's disease.

In a 12-week, double-blind, placebo-controlled trial, suicidal ideation was reported by 2% of patients treated with AUSTEDO, compared to no patients on placebo; no suicide attempts and no completed suicides were reported. Depression was reported by 4% of patients treated with AUSTEDO.

When considering the use of AUSTEDO, the risk of suicidality should be balanced against the need for treatment of chorea. All patients treated with AUSTEDO should be observed for new or worsening depression or suicidality. If depression or suicidality does not resolve, consider discontinuing treatment with AUSTEDO.

Patients, their caregivers, and families should be informed of the risks of depression, worsening depression, and suicidality associated with AUSTEDO, and should be instructed to report behaviors of concern promptly to the treating physician. Patients with Huntington's disease who express suicidal ideation should be evaluated immediately.

**5.3 Neuroleptic Malignant Syndrome (NMS)**

A potentially fatal symptom complex sometimes referred to as Neuroleptic Malignant Syndrome (NMS) has been reported in association with drugs that reduce dopaminergic transmission [see *Drug Interactions* (7.4)]. While NMS has not been observed in patients receiving AUSTEDO, it has been observed in patients receiving tetrabenazine (a closely related VMAT2 inhibitor). Clinicians should be alerted to the signs and symptoms associated with NMS. Clinical manifestations of NMS are hyperpyrexia, muscle rigidity, altered mental status, and evidence of autonomic instability (irregular pulse or blood pressure, tachycardia, diaphoresis, and cardiac dysrhythmia). Additional signs may include elevated creatinine phosphokinase, myoglobinuria, rhabdomyolysis, and acute renal failure. The diagnosis of NMS can be complicated; other serious medical illness (e.g., pneumonia, systemic infection) and untreated or inadequately treated extrapyramidal disorders can present with similar signs and symptoms. Other important considerations in the differential diagnosis include central anticholinergic toxicity, heat stroke, drug fever, and primary central nervous system pathology.

The management of NMS should include (1) immediate discontinuation of AUSTEDO; (2) intensive symptomatic treatment and medical monitoring; and (3) treatment of any concomitant serious medical problems for which specific treatments are available. There is no general agreement about specific pharmacological treatment regimens for NMS.

Recurrence of NMS has been reported with resumption of drug therapy. If treatment with AUSTEDO is needed after recovery from NMS, patients should be monitored for signs of recurrence.

**5.4 Akathisia, Agitation, and Restlessness**

AUSTEDO may increase the risk of akathisia, agitation, and restlessness in patients with Huntington's disease. In a 12-week, double-blind, placebo-controlled trial, akathisia, agitation, or restlessness was reported by 4% of patients treated with AUSTEDO, compared to 2% of patients on placebo.

Patients receiving AUSTEDO should be monitored for signs and symptoms of restlessness and agitation, as these may be indicators of developing akathisia. If a patient develops akathisia during treatment with AUSTEDO, the AUSTEDO dose should be reduced; some patients may require discontinuation of therapy.

**5.5 Parkinsonism**

AUSTEDO may cause parkinsonism in patients with Huntington's disease.

Because rigidity can develop as part of the underlying disease process in Huntington's disease, it may be difficult to distinguish between this potential drug-induced adverse reaction and progression of the underlying disease process. Drug-induced parkinsonism has the potential to cause more functional disability than untreated chorea for some patients with Huntington's disease. If a patient develops parkinsonism during treatment with AUSTEDO, the AUSTEDO dose should be reduced; some patients may require discontinuation of therapy.

**5.6 Sedation and Somnolence**

Sedation is a common dose-limiting adverse reaction of AUSTEDO. In a 12-week, double-blind, placebo-controlled trial, 11% of AUSTEDO-treated patients reported somnolence compared with 4% of patients on placebo and 9% of AUSTEDO-treated patients reported fatigue compared with 4% of placebo-treated patients.

Patients should not perform activities requiring mental alertness to maintain the safety of themselves or others, such as operating a motor vehicle or operating hazardous machinery, until they are on a maintenance dose of AUSTEDO and know how the drug affects them.

**5.7 QTc Prolongation**

Tetrabenazine, a closely related VMAT2 inhibitor, causes an increase (about 8 msec) in the corrected QT (QTc) interval [see *Clinical Pharmacology* (12.2)].

A clinically relevant QT prolongation may occur in some patients treated with AUSTEDO who are CYP2D6 poor metabolizers or are co-administered a strong CYP2D6 inhibitor [see *Clinical Pharmacology* (12.2, 12.3)].

The use of AUSTEDO should be avoided in combination with other drugs that are known to prolong QTc [see *Drug Interactions* (7.6)].

AUSTEDO should also be avoided in patients with congenital long QT syndrome and in patients with a history of cardiac arrhythmias. Certain circumstances may increase the risk of the occurrence of torsade de pointes and/or sudden death in association with the use of drugs that prolong the QTc interval, including (1) bradycardia; (2) hypokalemia or hypomagnesemia; (3) concomitant use of other drugs that prolong the QTc interval; and (4) presence of congenital prolongation of the QT interval.

**5.8 Hyperprolactinemia**

Serum prolactin levels were not evaluated in the AUSTEDO development program. Tetrabenazine, a closely related VMAT2 inhibitor, elevates serum prolactin concentrations in humans. Following administration of 25 mg of tetrabenazine to healthy volunteers, peak plasma prolactin levels increased 4- to 5-fold.

Tissue culture experiments indicate that approximately one-third of human breast cancers are prolactin-dependent *in vitro*, a factor of potential importance if AUSTEDO is being considered for a patient with previously detected breast cancer. Although amenorrhea, galactorrhea, gynecomastia, and impotence can be caused by elevated serum prolactin concentrations, the clinical significance of elevated serum prolactin concentrations for most patients is unknown.

Chronic increase in serum prolactin levels (although not evaluated in the AUSTEDO or tetrabenazine development programs) has been associated with low levels of estrogen and increased risk of osteoporosis. If there is a clinical suspicion of

symptomatic hyperprolactinemia, appropriate laboratory testing should be done and consideration should be given to discontinuation of AUSTEDO.

### 5.9 Binding to Melanin-Containing Tissues

Since deutetrabenazine or its metabolites bind to melanin-containing tissues, it could accumulate in these tissues over time. This raises the possibility that AUSTEDO may cause toxicity in these tissues after extended use. Neither ophthalmologic nor microscopic examination of the eye has been conducted in the chronic toxicity studies in a pigmented species such as dogs. Ophthalmologic monitoring in humans was inadequate to exclude the possibility of injury occurring after long-term exposure.

The clinical relevance of deutetrabenazine's binding to melanin-containing tissues is unknown. Although there are no specific recommendations for periodic ophthalmologic monitoring, prescribers should be aware of the possibility of long-term ophthalmologic effects [see *Clinical Pharmacology* (12.2)].

## 6 ADVERSE REACTIONS

The following serious adverse reactions are discussed in greater detail in other sections of the labeling:

- Depression and Suicidality [see *Warnings and Precautions* (5.2)]
- Neuroleptic Malignant Syndrome (NMS) [see *Warnings and Precautions* (5.3)]
- Akathisia, Agitation, and Restlessness [see *Warnings and Precautions* (5.4)]
- Parkinsonism [see *Warnings and Precautions* (5.5)]
- Sedation and Somnolence [see *Warnings and Precautions* (5.6)]
- QTc Prolongation [see *Warnings and Precautions* (5.7)]
- Hyperprolactinemia [see *Warnings and Precautions* (5.8)]
- Binding to Melanin-Containing Tissues [see *Warnings and Precautions* (5.9)]

### 6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Study 1 was a randomized, 12-week, placebo-controlled study in patients with chorea associated with Huntington's disease. A total of 45 patients received AUSTEDO, and 45 patients received placebo. Patients ranged in age between 23 and 74 years (mean 54 years); 56% were male, and 92% were Caucasian. The most common adverse reactions occurring in greater than 8% of AUSTEDO-treated patients were somnolence, diarrhea, dry mouth, and fatigue. Adverse reactions occurring in 4% or more of patients treated with AUSTEDO, and with a greater incidence than in patients on placebo, are summarized in Table 2.

**Table 2. Adverse Reactions in Study 1 Experienced by at Least 4% of Patients on AUSTEDO and with a Greater Incidence than on Placebo**

Adverse Reaction	AUSTEDO (N=45) %	Placebo (N=45) %
Somnolence	11	4
Diarrhea	9	0
Dry mouth	9	7
Fatigue	9	4
Urinary tract infection	7	2
Insomnia	7	4
Anxiety	4	2
Constipation	4	2
Contusion	4	2

One or more adverse reactions resulted in a reduction of the dose of study medication in 7% of patients in Study 1. The most common adverse reaction resulting in dose reduction in patients receiving AUSTEDO was dizziness (4%).

Agitation led to discontinuation in 2% of patients treated with AUSTEDO in Study 1.

## 7 DRUG INTERACTIONS

### 7.1 Strong CYP2D6 Inhibitors

A reduction in AUSTEDO dose may be necessary when adding a strong CYP2D6 inhibitor in patients maintained on a stable dose of AUSTEDO. Concomitant use of strong CYP2D6 inhibitors (e.g., paroxetine, fluoxetine, quinidine, bupropion) has been shown to increase the systemic exposure to the active dihydro-metabolites of deutetrabenazine by approximately 3-fold. The daily dose of AUSTEDO should not exceed 36 mg per day, and the maximum single dose of AUSTEDO should not exceed 18 mg in patients taking strong CYP2D6 inhibitors [see *Dosage and Administration* (2.3) and *Clinical Pharmacology* (12.3)].

### 7.2 Reserpine

Reserpine binds irreversibly to VMAT2 and the duration of its effect is several days. Prescribers should wait for chorea to reemerge before administering AUSTEDO to help reduce the risk of overdose and major depletion of serotonin and norepinephrine in the central nervous system. At least 20 days should elapse after stopping reserpine before starting AUSTEDO. AUSTEDO and reserpine should not be used concomitantly [see *Contraindications* (4)].

### 7.3 Monoamine Oxidase Inhibitors (MAOIs)

AUSTEDO is contraindicated in patients taking MAOIs. AUSTEDO should not be used in combination with an MAOI, or within 14 days of discontinuing therapy with an MAOI [see *Contraindications* (4)].

### 7.4 Neuroleptic Drugs

The risk for parkinsonism, NMS, and akathisia may be increased by concomitant use of AUSTEDO and dopamine antagonists or antipsychotics.

### 7.5 Alcohol or Other Sedating Drugs

Concomitant use of alcohol or other sedating drugs may have additive effects and worsen sedation and somnolence [see *Warnings and Precautions* (5.6)].

### 7.6 Drugs that Cause QTc Prolongation

Tetrabenazine, a closely related VMAT2 inhibitor, causes a small increase in the corrected QT (QTc) interval. Clinically relevant QT prolongation may also occur with AUSTEDO [see *Warnings and Precautions* (5.7), *Clinical Pharmacology* (12.2)].

The use of AUSTEDO should be avoided in combination with other drugs that are known to prolong QTc, including antipsychotic medications (e.g., chlorpromazine, haloperidol, thioridazine, ziprasidone), antibiotics (e.g., moxifloxacin), Class 1A (e.g., quinidine, procainamide), and Class III (e.g., amiodarone, sotalol) antiarrhythmic medications or any other medications known to prolong the QTc interval.

### 7.7 Tetrabenazine

AUSTEDO is contraindicated in patients currently taking tetrabenazine. AUSTEDO may be initiated the day following discontinuation of tetrabenazine [see *Dosage and Administration* (2.2)].

## 8 USE IN SPECIFIC POPULATIONS

### 8.1 Pregnancy

#### Risk Summary

There are no adequate data on the developmental risk associated with the use of AUSTEDO in pregnant women. Administration of deutetrabenazine to rats during organogenesis produced no clear adverse effect on embryofetal development. However, administration of tetrabenazine to rats throughout pregnancy and lactation resulted in an increase in stillbirths and postnatal offspring mortality [see *Data*].

In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2-4% and 15-20%, respectively. The background risk of major birth defects and miscarriage for the indicated population is unknown.

#### Data

##### Animal Data

Oral administration of deutetrabenazine (5, 10, or 30 mg/kg/day) or tetrabenazine (30 mg/kg/day) to pregnant rats during organogenesis had no clear effect on embryofetal development. The highest dose tested was 6 times the maximum recommended human dose of 48 mg/day, on a body surface area (mg/m<sup>2</sup>) basis.

The effects of deutetrabenazine when administered during organogenesis to rabbits or during pregnancy and lactation to rats have not been assessed.

Tetrabenazine had no effects on embryofetal development when administered to pregnant rabbits during the period of organogenesis at oral doses up to 60 mg/kg/day. When tetrabenazine was administered to female rats (doses of 5, 15, and 30 mg/kg/day) from the beginning of organogenesis through the lactation period, an increase in stillbirths and offspring postnatal mortality was observed at 15 and 30 mg/kg/day, and delayed pup maturation was observed at all doses.

### 8.2 Lactation

#### Risk Summary

There are no data on the presence of deutetrabenazine or its metabolites in human milk, the effects on the breastfed infant, or the effects of the drug on milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for AUSTEDO and any potential adverse effects on the breastfed infant from AUSTEDO or from the underlying maternal condition.

### 8.4 Pediatric Use

Safety and effectiveness in pediatric patients have not been established.

### 8.5 Geriatric Use

Clinical studies of AUSTEDO did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of hepatic, renal, and cardiac dysfunction, and of concomitant disease or other drug therapy.

### 8.6 Hepatic Impairment

The effect of hepatic impairment on the pharmacokinetics of deutetrabenazine and its primary metabolites has not been studied; however, in a clinical study conducted with tetrabenazine, a closely related VMAT2 inhibitor, there was a large increase in exposure to tetrabenazine and its active metabolites in patients with hepatic impairment. The clinical significance of this increased exposure has not been assessed, but because of concerns for a greater risk for serious adverse reactions, the use of AUSTEDO in patients with hepatic impairment is contraindicated [see *Contraindications* (4), *Clinical Pharmacology* (12.3)].

### 8.7 Poor CYP2D6 Metabolizers

Although the pharmacokinetics of deutetrabenazine and its metabolites have not been systematically evaluated in patients who do not express the drug metabolizing enzyme, it is likely that the exposure to  $\alpha$ -HTBZ and  $\beta$ -HTBZ would be increased similarly to taking a strong CYP2D6 inhibitor (approximately 3-fold). In patients who are CYP2D6 poor metabolizers, the daily dose of AUSTEDO should not exceed 36 mg (maximum single dose of 18 mg) [see *Dosage and Administration* (2.4) and *Clinical Pharmacology* (12.3)].


## 10 OVERDOSAGE

Overdoses ranging from 100 mg to 1 g have been reported in the literature with tetrabenazine, a closely related VMAT2 inhibitor. The following adverse reactions occurred with overdosing: acute dystonia, oculogyric crisis, nausea and vomiting, sweating, sedation, hypotension, confusion, diarrhea, hallucinations, rubor, and tremor.

Treatment should consist of those general measures employed in the management of overdose with any central nervous system-active drug. General supportive and symptomatic measures are recommended. Cardiac rhythm and vital signs should be monitored. In managing overdose, the possibility of multiple drug involvement should always be considered. The physician should consider contacting a poison control center on the treatment of any overdose. Telephone numbers for certified poison control centers are listed on the American Association of Poison Control Centers website [www.aapcc.org](http://www.aapcc.org).

## 11 DESCRIPTION

AUSTEDO (deutetrabenazine) is a vesicular monoamine transporter 2 (VMAT2) inhibitor for oral administration. The molecular weight of deutetrabenazine is 323.46; the pKa is 6.31. Deutetrabenazine is a hexahydro-dimethoxybenzoquinolizine derivative and has the following chemical name: (RR, SS)-1, 3, 4, 6, 7, 11b-hexahydro-9, 10-di(methoxy-d<sub>3</sub>)-3-(2-methylpropyl)-2H-benzo[a]quinolizine-2-one. The molecular formula for deutetrabenazine is C<sub>19</sub>H<sub>21</sub>D<sub>6</sub>NO<sub>3</sub>. Deutetrabenazine is a racemic mixture containing the following structures:



Deutetrabenazine is a white to slightly yellow crystalline powder that is sparingly soluble in water and soluble in ethanol.

AUSTEDO tablets contain 6 mg, 9 mg, or 12 mg deutetrabenazine, and the following inactive ingredients: ammonium hydroxide, black iron oxide, n-butyl alcohol, butylated hydroxyanisole, butylated hydroxytoluene, magnesium stearate, mannitol, microcrystalline cellulose, polyethylene glycol, polyethylene oxide, polysorbate 80, polyvinyl alcohol, povidone, propylene glycol, shellac, talc, titanium dioxide, and FD&C blue #2 lake. The 6 mg tablets also contain FD&C red #40 lake. The 12 mg tablets also contain FD&C yellow #6 lake.

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

The precise mechanism by which AUSTEDO (deutetrabenazine) exerts its anti-chorea effects is unknown but is believed to be related to its effect as a reversible depletor of monoamines (such as dopamine, serotonin, norepinephrine, and histamine) from nerve terminals. The major circulating metabolites ( $\alpha$ -dihydrodeutetrabenazine [HTBZ] and  $\beta$ -HTBZ) of deutetrabenazine, are reversible inhibitors of VMAT2, resulting in decreased uptake of monoamines into synaptic vesicles and depletion of monoamine stores.

### 12.2 Pharmacodynamics

#### Cardiac Electrophysiology

The effect of a single 12-mg or 24-mg dose of AUSTEDO on the QT interval was studied in a randomized, double-blind, placebo-controlled crossover study in healthy male and female subjects with moxifloxacin as a positive control. At 24 mg, AUSTEDO caused an approximately 4.5 msec mean increase in QTc (90% CI: 2.4, 6.5 msec). Effects at higher exposures to AUSTEDO or its metabolites have not been evaluated.

The effect of a single 25-mg or 50-mg dose of tetrabenazine, a closely related VMAT2 inhibitor, on the QT interval was studied in a randomized, double-blind, placebo-controlled crossover study in healthy male and female subjects with moxifloxacin as a positive control. At 50 mg, tetrabenazine caused an approximately 8 msec mean increase in QTc (90% CI: 5.0, 10.4 msec). Effects at higher exposures to either tetrabenazine or its metabolites have not been evaluated [see *Warnings and Precautions* (5.7) and *Drug Interactions* (7.6)].

#### Melanin Binding

Deutetrabenazine or its metabolites bind to melanin-containing tissues (i.e., eye, skin, fur) in pigmented rats. After a single oral dose of radiolabeled deutetrabenazine, radioactivity was still detected in eye and fur at 35 days following dosing [see *Warnings and Precautions* (5.9)].

### 12.3 Pharmacokinetics

After oral dosing up to 25 mg, plasma concentrations of deutetrabenazine are generally below the limit of detection because of the extensive hepatic metabolism of deutetrabenazine to the active deuterated dihydro metabolites (HTBZ),  $\alpha$ -HTBZ and  $\beta$ -HTBZ. Linear dose dependence of C<sub>max</sub> and AUC was observed for the active metabolites following single or multiple doses of deutetrabenazine (6 mg to 24 mg and 7.5 mg twice daily to 22.5 mg twice daily).

#### Absorption

Following oral administration of deutetrabenazine, the extent of absorption is at least 80%.

Plasma concentrations of deutetrabenazine are generally below the limit of detection after oral dosing. Peak plasma concentrations (C<sub>max</sub>) of deuterated  $\alpha$ -HTBZ and  $\beta$ -HTBZ are reached within 3 to 4 hours after dosing.

#### Effect of Food

The effects of food on the bioavailability of AUSTEDO were studied in subjects administered a single dose with and without food. Food had no effect on the area under the plasma concentration-time curve (AUC) of  $\alpha$ -HTBZ or  $\beta$ -HTBZ, although C<sub>max</sub> was increased by approximately 50% in the presence of food [see *Dosage and Administration* (2.1)].

#### Distribution

The median volume of distribution (V<sub>c</sub>/F) of the  $\alpha$ -HTBZ, and the  $\beta$ -HTBZ metabolites of AUSTEDO are approximately 500 L and 730 L, respectively.

Results of PET-scan studies in humans show that following intravenous injection of <sup>11</sup>C-labeled tetrabenazine or  $\alpha$ -HTBZ, radioactivity is rapidly distributed to the brain, with the highest binding in the striatum and lowest binding in the cortex.

The *in vitro* protein binding of tetrabenazine,  $\alpha$ -HTBZ, and  $\beta$ -HTBZ was examined in human plasma for concentrations ranging from 50 to 200 ng/mL. Tetrabenazine binding ranged from 82% to 85%,  $\alpha$ -HTBZ binding ranged from 60% to 68%, and  $\beta$ -HTBZ binding ranged from 59% to 63%.

#### Elimination

AUSTEDO is primarily renally eliminated in the form of metabolites.

The half-life of total ( $\alpha$ + $\beta$ )-HTBZ from deutetrabenazine is approximately 9 to 10 hours.

The median clearance values (CL/F) of the  $\alpha$ -HTBZ, and the  $\beta$ -HTBZ metabolites of AUSTEDO are approximately 47 L/hour and 70 L/hour, respectively, in the Huntington's disease patient population.

#### Metabolism

*In vitro* experiments in human liver microsomes demonstrate that deutetrabenazine is extensively biotransformed, mainly by carbonyl reductase, to its major active metabolites,  $\alpha$ -HTBZ and  $\beta$ -HTBZ, which are subsequently metabolized primarily by CYP2D6, with minor contributions of CYP1A2 and CYP3A4/5, to form several minor metabolites.

#### Excretion

In a mass balance study in 6 healthy subjects, 75% to 86% of the deutetrabenazine dose was excreted in the urine, and fecal recovery accounted for 8% to 11% of the dose. Urinary excretion of the  $\alpha$ -HTBZ and  $\beta$ -HTBZ metabolites from deutetrabenazine each accounted for less than 10% of the administered dose. Sulfate and glucuronide conjugates of the  $\alpha$ -HTBZ and  $\beta$ -HTBZ metabolites of deutetrabenazine, as well as products of oxidative metabolism, accounted for the majority of metabolites in the urine.

#### Specific Populations

##### Male and Female Patients

There is no apparent effect of gender on the pharmacokinetics of  $\alpha$ -HTBZ and  $\beta$ -HTBZ of deutetrabenazine.

##### Patients with Renal Impairment

No clinical studies have been conducted to assess the effect of renal impairment on the PK of AUSTEDO.

##### Patients with Hepatic Impairment

The effect of hepatic impairment on the pharmacokinetics of deutetrabenazine and its primary metabolites has not been studied. However, in a clinical study conducted to assess the effect of hepatic impairment on the pharmacokinetics of tetrabenazine, a closely related VMAT2 inhibitor, the exposure to  $\alpha$ -HTBZ and  $\beta$ -HTBZ was up to 40% greater in patients with hepatic impairment, and the mean tetrabenazine C<sub>max</sub> in patients with hepatic impairment was up to 190-fold higher than in healthy subjects [see *Contraindications* (4), *Use in Specific Populations* (8.6)].

##### Poor CYP2D6 Metabolizers

Although the pharmacokinetics of deutetrabenazine and its metabolites have not been systematically evaluated in patients who do not express the drug metabolizing enzyme CYP2D6, it is likely that the exposure to  $\alpha$ -HTBZ and  $\beta$ -HTBZ would be increased similarly to taking strong CYP2D6 inhibitors (approximately 3-fold) [see *Dosage and Administration* (2.4), *Drug Interactions* (7.1)].

#### Drug Interaction Studies

Deutetrabenazine,  $\alpha$ -HTBZ, and  $\beta$ -HTBZ have not been evaluated in *in vitro* studies for induction or inhibition of CYP enzymes or interaction with P-glycoprotein. The results of *in vitro* studies of tetrabenazine do not suggest that tetrabenazine or its  $\alpha$ -HTBZ or  $\beta$ -HTBZ metabolites are likely to result in clinically significant inhibition of CYP2D6, CYP1A2, CYP2B6, CYP2C8, CYP2C9, CYP2C19, CYP2E1, or CYP3A. *In vitro* studies suggest that neither tetrabenazine nor its  $\alpha$ -HTBZ or  $\beta$ -HTBZ metabolites are likely to result in clinically significant induction of CYP1A2, CYP3A4, CYP2B6, CYP2C8, CYP2C9, or CYP2C19. Neither tetrabenazine nor its  $\alpha$ -HTBZ or  $\beta$ -HTBZ metabolites are likely to be a substrate or inhibitor of P-glycoprotein at clinically relevant concentrations *in vivo*.

The deutetrabenazine metabolites, 2-methylpropanoic acid of  $\beta$ -HTBZ (M1) and monohydroxy tetrabenazine (M4), have been evaluated in a panel of *in vitro* drug-drug interaction studies; the results indicate that M1/M4 are not expected to cause clinically relevant drug interactions.

#### CYP2D6 Inhibitors

*In vitro* studies indicate that the  $\alpha$ -HTBZ and  $\beta$ -HTBZ metabolites of deutetrabenazine are substrates for CYP2D6. The effect of CYP2D6 inhibition on the pharmacokinetics of deutetrabenazine and its metabolites was studied in 24 healthy subjects following a single 22.5 mg dose of deutetrabenazine given after 8 days of administration of the strong CYP2D6 inhibitor paroxetine 20 mg daily. In the presence of paroxetine, systemic exposure (AUC<sub>inf</sub>) of  $\alpha$ -HTBZ was 1.9-fold higher and  $\beta$ -HTBZ was 6.5-fold higher, resulting in approximately 3-fold increase in AUC<sub>inf</sub> for total ( $\alpha$ + $\beta$ )-HTBZ. Paroxetine decreased the clearance of  $\alpha$ -HTBZ and  $\beta$ -HTBZ metabolites of AUSTEDO with corresponding increases in mean half-life of approximately 1.5-fold and 2.7-fold, respectively. In the presence of paroxetine, C<sub>max</sub> of  $\alpha$ -HTBZ and  $\beta$ -HTBZ were 1.2-fold and 2.2-fold higher, respectively.

The effect of moderate or weak CYP2D6 inhibitors such as duloxetine, terbinafine, amiodarone, or sertraline on the exposure of deutetrabenazine and its metabolites has not been evaluated.

**Digoxin**

AUSTEDO was not evaluated for interaction with digoxin. Digoxin is a substrate for P-glycoprotein. A study in healthy subjects showed that tetrabenazine (25 mg twice daily for 3 days) did not affect the bioavailability of digoxin, suggesting that at this dose, tetrabenazine does not affect P-glycoprotein in the intestinal tract. *In vitro* studies also do not suggest that tetrabenazine or its metabolites are P-glycoprotein inhibitors.

**13 NONCLINICAL TOXICOLOGY****13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility****Carcinogenesis**

No carcinogenicity studies were performed with deutetrabenazine.

No increase in tumors was observed in p53<sup>+/−</sup> transgenic mice treated orally with tetrabenazine at doses of 0, 5, 15, and 30 mg/kg/day for 26 weeks.

**Mutagenesis**

Deutetrabenazine and its deuterated  $\alpha$ -HTBZ and  $\beta$ -HTBZ metabolites were negative in *in vitro* (bacterial reverse mutation and chromosome aberration in human peripheral blood lymphocytes) assays in the presence or absence of metabolic activation and in the *in vivo* micronucleus assay in mice.

**Impairment of Fertility**

The effects of deutetrabenazine on fertility have not been evaluated. Oral administration of deutetrabenazine (doses of 5, 10, or 30 mg/kg/day) to female rats for 3 months resulted in estrous cycle disruption at all doses; the lowest dose tested was similar to the maximum recommended human dose (48 mg/day) on a body surface area (mg/m<sup>2</sup>) basis.

Oral administration of tetrabenazine (doses of 5, 15, or 30 mg/kg/day) to female rats prior to and throughout mating, and continuing through day 7 of gestation, resulted in disrupted estrous cyclicity at doses greater than 5 mg/kg/day. No effects on mating and fertility indices or sperm parameters (motility, count, density) were observed when males were treated orally with tetrabenazine at doses of 5, 15 or 30 mg/kg/day prior to and throughout mating with untreated females.

**14 CLINICAL STUDIES****Double-Blind, Placebo-Controlled Study**

The efficacy of AUSTEDO as a treatment for chorea associated with Huntington's disease was established primarily in Study 1, a randomized, double-blind, placebo-controlled, multi-center trial conducted in 90 ambulatory patients with manifest chorea associated with Huntington's disease. The diagnosis of Huntington's disease was based on family history, neurological exam, and genetic testing. Treatment duration was 12 weeks, including an 8-week dose titration period and a 4-week maintenance period, followed by a 1-week washout. Patients were not blinded to discontinuation. AUSTEDO was started at 6 mg per day and titrated upward, at weekly intervals, in 6 mg increments until satisfactory treatment of chorea was achieved, intolerable side effects occurred, or until a maximal dose of 48 mg per day was reached. The primary efficacy endpoint was the Total Maximal Chorea Score, an item of the Unified Huntington's Disease Rating Scale (UHDRS). On this scale, chorea is rated from 0 to 4 (with 0 representing no chorea) for 7 different parts of the body. The total score ranges from 0 to 28.

Of the 90 patients enrolled, 87 patients completed the study. The mean age was 54 (range 23 to 74). Patients were 56% male and 92% Caucasian. The mean dose after titration was 40 mg per day. Table 3 and Figure 1 summarize the effects of AUSTEDO on chorea based on the Total Maximal Chorea Score. Total Maximal Chorea Scores for patients receiving AUSTEDO improved by approximately 4.4 units from baseline to the maintenance period (average of Week 9 and Week 12), compared to approximately 1.9 units in the placebo group. The treatment effect of -2.5 units was statistically significant ( $p < 0.0001$ ). The Maintenance Endpoint is the mean of the Total Maximal Chorea Scores for the Week 9 and Week 12 visits. At the Week 13 follow-up visit (1 week after discontinuation of the study medication), the Total Maximal Chorea Scores of patients who had received AUSTEDO returned to baseline (Figure 1).


**Table 3. Change from Baseline to Maintenance Therapy in Total Maximal Chorea (TMC)<sup>a</sup> Score in Patients Treated with AUSTEDO in Study 1**

Motor Endpoint	AUSTEDO N = 45	Placebo N = 45	p value
Change in Total Chorea Score <sup>a</sup> from Baseline to Maintenance Therapy <sup>b</sup>	-4.4	-1.9	<0.0001

<sup>a</sup>TMC is a subscale of the Unified Huntington's Disease Rating Scale (UHDRS)

<sup>b</sup>Primary efficacy endpoint

**Figure 1. Total Maximal Chorea Score Over Time in Study 1**



**Figure 2. Distribution of the Change in Total Maximal Chorea Scores in Study 1**




Figure 2 shows the distribution of values for the change in Total Maximal Chorea Score in Study 1. Negative values indicate a reduction in chorea and positive numbers indicate an increase in chorea.

A patient-rated global impression of change assessed how patients rated their overall Huntington's disease symptoms. Fifty-one percent of patients treated with AUSTEDO rated their symptoms as "Much Improved" or "Very Much Improved" at the end of treatment, compared to 20% of placebo-treated patients.

In a physician-rated clinical global impression of change, 42% percent of patients treated with AUSTEDO rated their symptoms as "Much Improved" or "Very Much Improved" at the end of treatment compared to 13% of placebo-treated patients.

**16 HOW SUPPLIED/STORAGE AND HANDLING****16.1 How Supplied**

AUSTEDO tablets are available in the following strengths and packages:

6 mg: round, purple-coated tablets, with "SD" over "6" printed in black ink on one side.

Bottles of 60 tablets: NDC 68546-170-60.

9 mg: round, blue-coated tablets, with "SD" over "9" printed in black ink on one side.

Bottles of 60 tablets: NDC 68546-171-60.

12 mg: round, beige-coated tablets, with "SD" over "12" printed in black ink on one side.

Bottles of 60 tablets: NDC 68546-172-60.

**16.2 Storage**

Store at 25° C (77°F); excursions permitted to 15° C to 30° C (59° F to 86° F) [see USP Controlled Room Temperature]. Protect from light and moisture.

**17 PATIENT COUNSELING INFORMATION**

Advise the patient or caregiver to read the FDA-approved patient labeling (Medication Guide).

**Administration Instructions**

Advise patients to take AUSTEDO with food. AUSTEDO tablets should be swallowed whole and not chewed, crushed, or broken [see Dosage and Administration (2.1)].

**Risk of Depression and Suicide**

Advise patients, their caregivers, and families that AUSTEDO may increase the risk of depression, worsening depression, and suicidality, and to immediately report any symptoms to a healthcare provider [see Contraindications (4), Warnings and Precautions (5.2)].

**Risk of Sedation and Somnolence**

Advise patients that AUSTEDO may cause sedation and somnolence and may impair the ability to perform tasks that require complex motor and mental skills. Until they learn how they respond to a stable dose of AUSTEDO, patients should be careful doing activities that require them to be alert, such as driving a car or operating machinery [see *Warnings and Precautions (5.6)*].

**Interaction with Alcohol or Other Sedating Drugs**

Advise patients that alcohol or other drugs that cause sleepiness will worsen somnolence [see *Drug Interactions (7.5)*].

**Concomitant Medications**

Advise patients to notify their physician of all medications they are taking and to consult with their healthcare provider before starting any new medications because of a potential for interactions [see *Contraindications (4)* and *Drug Interactions (7.1, 7.4, 7.5, 7.6)*].

Distributed by:

**Teva Pharmaceuticals USA, Inc.**

North Wales, PA 19454

AUSTEDO™ is a trademark of Auspex Pharmaceuticals Inc., a member of the Teva Group.

AUS-001

U.S. Patent No: 8,524,733

**MEDICATION GUIDE**  
**AUSTEDO™ (aw-*STED-oh*)**  
**(deutetrabenazine)**  
**tablets, for oral use**

**What is the most important information I should know about AUSTEDO?**

- AUSTEDO can cause serious side effects, including:
  - **depression**
  - **suicidal thoughts**
  - **suicidal actions**
- **Do not** start taking AUSTEDO if you are depressed (have untreated depression or depression that is not well controlled by medicine) **or** have suicidal thoughts.
- Pay close attention to any changes, especially sudden changes, in mood, behaviors, thoughts or feelings. This is especially important when AUSTEDO is started and when the dose is changed.

**Call your healthcare provider right away if you become depressed or have any of the following symptoms, especially if they are new, worse, or worry you:**

- feel sad or have crying spells
- lose interest in seeing your friends or doing things you used to enjoy
- sleep a lot **more** or a lot **less** than usual
- feel unimportant
- feel guilty
- feel hopeless or helpless
- feel more irritable, angry, or aggressive than usual
- feel more or less hungry than usual or notice a big change in your body weight
- have trouble paying attention
- feel tired or sleepy all the time
- have thoughts about hurting yourself or ending your life

**What is AUSTEDO?**

AUSTEDO is a prescription medicine that is used to treat the involuntary movements (chorea) of Huntington's disease. AUSTEDO does not cure the cause of the involuntary movements, and it does not treat other symptoms of Huntington's disease, such as problems with thinking or emotions.

It is not known if AUSTEDO is safe and effective in children.

**Who should not take AUSTEDO?****Do not take AUSTEDO if you:**

- are depressed or have thoughts of suicide. See "**What is the most important information I should know about AUSTEDO?**"
- have liver problems.
- are taking a monoamine oxidase inhibitor (MAOI) medicine. **Do not** take an MAOI within 14 days after you stop taking AUSTEDO. **Do not** start AUSTEDO if you stopped taking an MAOI in the last 14 days. Ask your healthcare provider or pharmacist if you are not sure.
- are taking reserpine. **Do not** take medicines that contain reserpine (such as Serpalan® and Renese®-R) with AUSTEDO. If your healthcare provider plans to switch you from taking reserpine to AUSTEDO, you must wait at least 20 days after your last dose of reserpine before you start taking AUSTEDO.
- **are taking tetrabenazine (XENAZINE®). If your healthcare provider plans to switch you from tetrabenazine (XENAZINE®) to AUSTEDO, take your first dose of AUSTEDO on the day after your last dose of tetrabenazine (XENAZINE®).**

**Before taking AUSTEDO, tell your healthcare provider about all of your medical conditions, including if you:**

- have emotional or mental problems (for example, depression, nervousness, anxiety, anger, agitation, psychosis, previous suicidal thoughts or suicide attempts).
- have liver disease.
- have an irregular heartbeat (cardiac arrhythmia) or a heart problem called congenital long QT syndrome.
- have breast cancer or a history of breast cancer.
- are pregnant or plan to become pregnant. It is not known if AUSTEDO can harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if AUSTEDO passes into breast milk.

**Tell your healthcare provider about all of the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Using AUSTEDO with certain other medicines may cause side effects. Do not start any new medicines while taking AUSTEDO without talking to your healthcare provider first.****How should I take AUSTEDO?**

- Take AUSTEDO by mouth.
- Take AUSTEDO exactly as your healthcare provider tells you to take it.
- Take AUSTEDO with food. Swallow AUSTEDO tablets whole with water. Do not chew, crush, or break AUSTEDO tablets before swallowing. If you cannot swallow AUSTEDO tablets whole, tell your healthcare provider. You may need a different medicine.
- If your dose of AUSTEDO is 12 mg or more each day, take AUSTEDO tablets 2 times a day in equal doses with food.
- Your healthcare provider will increase your dose of AUSTEDO each week for several weeks, until you and your healthcare provider find the right dose for you.
- Tell your healthcare provider if you stop taking AUSTEDO for more than **1 week. Do not take another dose until you talk to your healthcare provider.**

**What should I avoid while taking AUSTEDO?**

Sleepiness (sedation) is a common side effect of AUSTEDO. While taking AUSTEDO, **do not drive a car or operate dangerous machinery until you know how AUSTEDO affects you.** Drinking alcohol and taking other drugs that may also cause sleepiness while you are taking AUSTEDO may increase any sleepiness caused by AUSTEDO.

**What are the possible side effects of AUSTEDO?**

**AUSTEDO can cause serious side effects, including:**

- **Depression and suicidal thoughts or actions.** See “What is the most important information I should know about AUSTEDO?”
- **Neuroleptic Malignant Syndrome (NMS).** Call your healthcare provider right away and go to the nearest emergency room if you develop these signs and symptoms that do not have another obvious cause:
  - high fever
  - problems thinking
  - increased sweating
  - stiff muscles
  - very fast or uneven heartbeat
- **Restlessness.** You may get a condition where you feel a strong urge to move. This is called akathisia.
- **Parkinsonism.** Symptoms of parkinsonism include: slight shaking, body stiffness, trouble moving or keeping your balance.
- **Irregular heartbeat.** AUSTEDO increases your chance of having certain changes in the electrical activity in your heart. These changes can lead to a dangerous abnormal heartbeat. Taking AUSTEDO with certain medicines may increase this chance.

**The most common side effects of AUSTEDO include:**

- sleepiness (sedation)
- diarrhea
- tiredness
- dry mouth

These are not all the possible side effects of AUSTEDO. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**How should I store AUSTEDO?**

- Store AUSTEDO tablets at room temperature, between 68°F to 77°F (20°C to 25°C).
- Keep the bottle tightly closed to protect AUSTEDO from light and moisture.

**Keep AUSTEDO tablets and all medications out of reach of children.**

**General information about the safe and effective use of AUSTEDO.**

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use AUSTEDO for a condition for which it was not prescribed. Do not give AUSTEDO to other people, even if they have the same symptoms you have. It may harm them. You can ask your pharmacist or healthcare provider for information about AUSTEDO that is written for health professionals.

**What are the ingredients in AUSTEDO?**

**Active ingredient:** deutetrabenazine

**Inactive ingredients:** ammonium hydroxide, black iron oxide, n-butyl alcohol, butylated hydroxyanisole, butylated hydroxytoluene, magnesium stearate, mannitol, microcrystalline cellulose, polyethylene glycol, polyethylene oxide, polysorbate 80, polyvinyl alcohol, povidone, propylene glycol, shellac, talc, titanium dioxide and FD&C blue #2 lake. The 6 mg tablets also contain FD&C red #40 lake. The 12 mg tablets also contain FD&C yellow #6 lake.

Distributed by:

**Teva Pharmaceuticals USA, Inc.**

North Wales, PA 19454

AUSTEDO™ is a trademark of Auspex Pharmaceuticals Inc., a member of the Teva Group.

AUSMG-001

For more information, go to [www.AUSTEDO.com](http://www.AUSTEDO.com) or call 1-888-483-8279.

This Medication Guide has been approved by the U.S. Food and Drug Administration. Issued: 4/2017