



**Reach out to us, we'll  
reach for solutions together**

Teva **Shared Solutions**<sup>®</sup> is your partner from the very beginning of treatment, from starting your prescription to financial assistance. Let the broad reach of our services guide you toward a plan for starting and reaching your treatment goals with AUSTEDO<sup>®</sup> XR (deutetrabenazine) extended-release tablets.

## Sign up for Shared Solutions

by phone Monday-Friday  
from 8AM to 8PM CT at  
**1-800-887-8100**

or

through our  
website anytime at  
**MySharedSolutions.com**

### APPROVED USES

**AUSTEDO<sup>®</sup> XR (deutetrabenazine) extended-release tablets and AUSTEDO<sup>®</sup> (deutetrabenazine) tablets are prescription medicines that are used to treat:**

- the involuntary movements (chorea) of Huntington's disease. AUSTEDO XR and AUSTEDO do not cure the cause of the involuntary movements, and it does not treat other symptoms of Huntington's disease, such as problems with thinking or emotions.
- movements in the face, tongue, or other body parts that cannot be controlled (tardive dyskinesia).




It is not known if AUSTEDO XR and AUSTEDO are safe and effective in children.

### IMPORTANT SAFETY INFORMATION

**AUSTEDO XR and AUSTEDO can cause serious side effects in people with Huntington's disease, including: depression, suicidal thoughts, or suicidal actions. Do not** start taking AUSTEDO XR or AUSTEDO if you are depressed (have untreated depression or depression that is not well controlled by medicine) **or** have suicidal thoughts. Pay close attention to any changes, especially sudden changes, in mood, behaviors, thoughts or feelings. This is especially important when AUSTEDO XR or AUSTEDO is started and when the dose is changed. Call your healthcare provider right away if you become depressed, have unusual changes in mood or behavior, or have thoughts of suicide.

**Please read Important Safety Information on pages 6-7 and the accompanying Medication Guide.**

## What is Teva **Shared Solutions**<sup>®</sup>, and how can it help me?

With patient support from **Shared Solutions**, you'll have a Patient Access Specialist for  insurance and  pharmacy questions and  Nurse support for treatment and dosing guidance. And because your **Shared Solutions** team are dedicated professionals in their field, you'll get real help from real people.



## Can Teva **Shared Solutions**<sup>®</sup> help make my prescription affordable?

**Shared Solutions** helps with financial, insurance, and prescription assistance so you can start treatment sooner and stay on treatment. Whether you're insured, underinsured, or not insured at all, we'll verify coverage and work through financial assistance options to help make treatment affordable. Plus, we'll help you navigate the paperwork, so nothing falls through the cracks. **Close to 90% of existing patients on AUSTEDO (deutetrabenazine) tablets pay \$10 or less per month** for their prescription. We'll help you explore your financial options.



Start AUSTEDO with us, we'll help you find your financial path forward. **You may pay as little as:**

ELIGIBLE  
PATIENTS MAY  
PAY AS LITTLE AS

**\$0** **COPAY**  
PER MONTH\*

\*Offer is only available to patients with commercial insurance. Offer is NOT available to patients eligible for Medicare, Medicaid, or any other government-payer coverage. Out-of-pocket costs may vary. Exclusions and limitations apply. Please see complete Terms and Conditions at [AUSTEDO.com](https://www.austedo.com).



## Will Teva **Shared Solutions**<sup>®</sup> work with my pharmacy?

When your doctor decides AUSTEDO<sup>®</sup> XR (deutetrabenazine) extended-release tablets is right for you, you get your prescription in one of two ways:


### **Without Shared Solutions:**

Your doctor writes you a prescription. Your pharmacy (or sometimes a specialty pharmacy) fills it for you. Navigating insurance, copays, and financial questions is up to you.

or

### **With Shared Solutions:**

Your doctor sends your prescription and enrollment form to **Shared Solutions**, and we help you work through the details.



### ***What is a Specialty Pharmacy?***


A specialty pharmacy dispenses medications that require special handling, like temperature control, personalized monitoring, or frequent dosage adjustments.



## How can Teva **Shared Solutions®** help once I'm on treatment?

While taking AUSTEDO® XR (deutetrabenzine) extended-release tablets with support from **Shared Solutions**, you will receive calls throughout treatment from our Nurses, who can help you with things like:

- ✓ Learning about your diagnosis and AUSTEDO XR
- ✓ Starting a treatment plan, dosing guidance, and even managing your refills, to keep you on track and on treatment
- ✓ How to take and how to remember to take your AUSTEDO XR
- ✓ Providing tools and resources to help manage your treatment



*It can be overwhelming to figure out how a new treatment works; that's why we set up a program to support our patients every step of the way.*

- **Shared Solutions** Nurse

## APPROVED USES

**AUSTEDO® XR (deutetrabenazine) extended-release tablets and AUSTEDO® (deutetrabenazine) tablets are prescription medicines that are used to treat:**

- the involuntary movements (chorea) of Huntington's disease. AUSTEDO XR and AUSTEDO do not cure the cause of the involuntary movements, and it does not treat other symptoms of Huntington's disease, such as problems with thinking or emotions.
- movements in the face, tongue, or other body parts that cannot be controlled (tardive dyskinesia).

It is not known if AUSTEDO XR and AUSTEDO are safe and effective in children.

## IMPORTANT SAFETY INFORMATION

**AUSTEDO XR and AUSTEDO can cause serious side effects in people with Huntington's disease, including: depression, suicidal thoughts, or suicidal actions.**

**Do not** start taking AUSTEDO XR or AUSTEDO if you are depressed (have untreated depression or depression that is not well controlled by medicine) **or** have suicidal thoughts. Pay close attention to any changes, especially sudden changes, in mood, behaviors, thoughts or feelings. This is especially important when AUSTEDO XR or AUSTEDO is started and when the dose is changed. Call your healthcare provider right away if you become depressed, have unusual changes in mood or behavior, or have thoughts of suicide.

**Do not take AUSTEDO XR or AUSTEDO if you:**

- have Huntington's disease and are depressed or have thoughts of suicide.
- have liver problems.
- are taking reserpine. **Do not** take medicines that contain reserpine with AUSTEDO XR or AUSTEDO. If your healthcare provider plans to switch you from taking reserpine to AUSTEDO XR or AUSTEDO, you must wait at least 20 days after your last dose of reserpine before you start taking AUSTEDO XR or AUSTEDO.
- are taking a monoamine oxidase inhibitor (MAOI) medicine. **Do not** take an MAOI within 14 days after you stop taking AUSTEDO XR or AUSTEDO. **Do not** start AUSTEDO XR or AUSTEDO if you stopped taking an MAOI in the last 14 days. Ask your healthcare provider or pharmacist if you are not sure.
- are taking tetrabenazine. If your healthcare provider plans to switch you from tetrabenazine to AUSTEDO XR or AUSTEDO, take your first dose of AUSTEDO XR or AUSTEDO on the day after your last dose of tetrabenazine.
- are taking valbenazine.

## IMPORTANT SAFETY INFORMATION (CONTINUED)

### Other possible serious side effects include:

- **Irregular heartbeat (QT prolongation).** AUSTEDO XR and AUSTEDO increases your chance of having certain changes in the electrical activity in your heart. These changes can lead to a dangerous abnormal heartbeat. Taking AUSTEDO XR or AUSTEDO with certain medicines may increase this chance.
- **Neuroleptic Malignant Syndrome.** Call your healthcare provider right away and go to the nearest emergency room if you develop these signs and symptoms that do not have another obvious cause: high fever, stiff muscles, problems thinking, very fast or uneven heartbeat, or increased sweating.
- **Restlessness.** You may get a condition where you feel a strong urge to move. This is called akathisia.
- **Parkinsonism.** Symptoms include: slight shaking, body stiffness, trouble moving, trouble keeping your balance, or falls.

### Sleepiness (sedation) is a common side effect of AUSTEDO XR and AUSTEDO.

While taking AUSTEDO XR or AUSTEDO, do not drive a car or operate dangerous machinery until you know how AUSTEDO XR or AUSTEDO affects you. Drinking alcohol and taking other drugs that may also cause sleepiness while you are taking AUSTEDO XR or AUSTEDO may increase any sleepiness caused by AUSTEDO XR and AUSTEDO.

**The most common side effects of AUSTEDO in people with Huntington's disease include** sleepiness (sedation), diarrhea, tiredness, and dry mouth.

**The most common side effects of AUSTEDO in people with tardive dyskinesia include** inflammation of the nose and throat (nasopharyngitis) and problems sleeping (insomnia).

The most common side effects of AUSTEDO XR are expected to be similar to AUSTEDO in people with Huntington's disease or tardive dyskinesia.

These are not all the possible side effects of AUSTEDO XR or AUSTEDO. Call your doctor for medical advice about side effects. You are encouraged to report side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch) or call 1-800-FDA-1088.

Please read the accompanying [Medication Guide](#).

© 2023 Teva Neuroscience, Inc. | AUS-45068 March 2023

## How do I sign up for Teva Shared Solutions<sup>®</sup> ?

Take advantage of support services, including options for financial assistance, nurse support, and benefits coverage.

### Sign up for Shared Solutions

by phone Monday-Friday  
from 8AM to 8PM CT at  
**1-800-887-8100**

or

through our  
website anytime at  
**MySharedSolutions.com**



**MEDICATION GUIDE**  
**AUSTEDO<sup>®</sup> XR (*aw-STED-oh XR*)**  
**(deutetrabenazine) extended-release tablets, for oral use**  
**AUSTEDO<sup>®</sup> (*aw-STED-oh*)**  
**(deutetrabenazine) tablets, for oral use**

**What is the most important information I should know about AUSTEDO XR and AUSTEDO?**

- AUSTEDO XR and AUSTEDO can cause serious side effects in people with Huntington's disease, including:
  - depression
  - suicidal thoughts
  - suicidal actions
- Do not start taking AUSTEDO XR or AUSTEDO if you have Huntington's disease and are depressed (have untreated depression or depression that is not well controlled by medicine) or have suicidal thoughts.
- Pay close attention to any changes, especially sudden changes, in mood, behaviors, thoughts, or feelings. This is especially important when AUSTEDO XR or AUSTEDO is started and when the dose is changed.

**Call your healthcare provider right away if you become depressed or have any of the following symptoms, especially if they are new, worse, or worry you:**

- feel sad or have crying spells
- lose interest in seeing your friends or doing things you used to enjoy
- sleep a lot more or a lot less than usual
- feel unimportant
- feel guilty
- feel hopeless or helpless
- feel more irritable, angry, or aggressive than usual
- feel more or less hungry than usual or notice a big change in your body weight
- have trouble paying attention
- feel tired or sleepy all the time
- have thoughts about hurting yourself or ending your life

**What are AUSTEDO XR and AUSTEDO?**

AUSTEDO XR and AUSTEDO are prescription medicines that are used to treat:

- the involuntary movements (chorea) of Huntington's disease. AUSTEDO XR and AUSTEDO do not cure the cause of the involuntary movements, and it does not treat other symptoms of Huntington's disease, such as problems with thinking or emotions.
- movements in the face, tongue, or other body parts that cannot be controlled (tardive dyskinesia).

It is not known if AUSTEDO XR and AUSTEDO are safe and effective in children.

**Who should not take AUSTEDO XR or AUSTEDO?**

**Do not take AUSTEDO XR or AUSTEDO if you:**

- have Huntington's disease and are depressed or have thoughts of suicide. See "What is the most important information I should know about AUSTEDO XR and AUSTEDO?"
- have liver problems.
- are taking reserpine. **Do not** take medicines that contain reserpine with AUSTEDO XR or AUSTEDO. If your healthcare provider plans to switch you from taking reserpine to AUSTEDO XR or AUSTEDO, you must wait at least 20 days after your last dose of reserpine before you start taking AUSTEDO XR or AUSTEDO.
- are taking a monoamine oxidase inhibitor (MAOI) medicine. **Do not** take an MAOI within 14 days after you stop taking AUSTEDO XR or AUSTEDO. **Do not** start AUSTEDO XR or AUSTEDO if you stopped taking an MAOI in the last 14 days. Ask your healthcare provider or pharmacist if you are not sure.
- are taking tetrabenazine. If your healthcare provider plans to switch you from tetrabenazine to AUSTEDO XR or AUSTEDO, take your first dose of AUSTEDO XR or AUSTEDO on the day after your last dose of tetrabenazine.
- are taking valbenazine.

**Before taking AUSTEDO XR or AUSTEDO, tell your healthcare provider about all of your medical conditions, including if you:**

- have emotional or mental problems (for example, depression, nervousness, anxiety, anger, agitation, psychosis, previous suicidal thoughts or suicide attempts).
- have liver disease.
- have an irregular heart rhythm or heartbeat (QT prolongation, cardiac arrhythmia) or a heart problem called congenital long QT syndrome.
- have low levels of potassium or magnesium in your blood (hypokalemia or hypomagnesemia).
- have breast cancer or a history of breast cancer.
- are pregnant or plan to become pregnant. It is not known if AUSTEDO XR or AUSTEDO can harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if AUSTEDO XR or AUSTEDO passes into breast milk.

**Tell your healthcare provider about all of the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Taking AUSTEDO XR or AUSTEDO with certain other medicines may cause side effects. Do not start any new medicines while taking AUSTEDO XR or AUSTEDO without talking to your healthcare provider first.

**How should I take AUSTEDO XR or AUSTEDO?**

- Take AUSTEDO XR or AUSTEDO exactly as your healthcare provider tells you to take it.
- If you take AUSTEDO XR, take your dose by mouth one time each day, with or without food.
- If you take AUSTEDO, take your dose by mouth and with food. If your dose of AUSTEDO is 12 mg or more each day, take AUSTEDO tablets 2 times a day in equal doses.
- Swallow AUSTEDO XR or AUSTEDO tablets whole with water. Do not chew, crush, or break AUSTEDO XR or AUSTEDO tablets before swallowing. If you cannot swallow AUSTEDO XR or AUSTEDO tablets whole, tell your healthcare provider. You may need a different medicine.
- Your healthcare provider may increase your dose of AUSTEDO XR or AUSTEDO each week for several weeks, until you and your healthcare provider find the right dose for you.
- Tell your healthcare provider if you stop taking AUSTEDO XR or AUSTEDO for more than 1 week. Do not take another dose until you talk to your healthcare provider.

*continued*

**What should I avoid while taking AUSTEDO XR or AUSTEDO?**

Sleepiness (sedation) is a common side effect of AUSTEDO XR and AUSTEDO. While taking AUSTEDO XR or AUSTEDO, do not drive a car or operate dangerous machinery until you know how AUSTEDO XR or AUSTEDO affects you. Drinking alcohol and taking other drugs that may also cause sleepiness while you are taking AUSTEDO XR or AUSTEDO may increase any sleepiness caused by AUSTEDO XR and AUSTEDO.

**What are the possible side effects of AUSTEDO XR and AUSTEDO?**

AUSTEDO XR and AUSTEDO can cause serious side effects, including:

- **Depression and suicidal thoughts or actions in people with Huntington's disease.** See "What is the most important information I should know about AUSTEDO XR and AUSTEDO?"
- **Irregular heartbeat (QT prolongation).** AUSTEDO XR and AUSTEDO increases your chance of having certain changes in the electrical activity in your heart. These changes can lead to a dangerous abnormal heartbeat. Taking AUSTEDO XR or AUSTEDO with certain medicines may increase this chance.
- **Neuroleptic Malignant Syndrome (NMS).** Call your healthcare provider right away and go to the nearest emergency room if you develop these signs and symptoms that do not have another obvious cause:
  - high fever
  - stiff muscles
  - problems thinking
  - very fast or uneven heartbeat
  - increased sweating
- **Restlessness.** You may get a condition where you feel a strong urge to move. This is called akathisia.
- **Parkinsonism.** Symptoms of parkinsonism include: slight shaking, body stiffness, trouble moving, trouble keeping your balance, or falls.

**The most common side effects of AUSTEDO in people with Huntington's disease include:**

- sleepiness (sedation)
- tiredness
- diarrhea
- dry mouth

**The most common side effects of AUSTEDO in people with tardive dyskinesia include:**

- inflammation of the nose and throat (nasopharyngitis)
- problems sleeping (insomnia)

The most common side effects of AUSTEDO XR are expected to be similar to AUSTEDO in people with Huntington's disease or tardive dyskinesia.

These are not all the possible side effects of AUSTEDO XR or AUSTEDO. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**How should I store AUSTEDO XR and AUSTEDO?**

- Store AUSTEDO XR and AUSTEDO tablets at room temperature between 68°F to 77°F (20°C to 25°C).
- Keep the bottle tightly closed to protect AUSTEDO XR and AUSTEDO from light and moisture.
- Do not throw away the desiccant canister in the bottle until the last dose of AUSTEDO XR or AUSTEDO is taken.

**Keep AUSTEDO XR and AUSTEDO and all medicines out of the reach of children.**

**General information about the safe and effective use of AUSTEDO XR and AUSTEDO.**

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use AUSTEDO XR or AUSTEDO for a condition for which it was not prescribed. Do not give AUSTEDO XR or AUSTEDO to other people, even if they have the same symptoms you have. It may harm them. You can ask your pharmacist or healthcare provider for information about AUSTEDO XR and AUSTEDO that is written for health professionals.

**What are the ingredients in AUSTEDO XR and AUSTEDO?****AUSTEDO XR:**

**Active ingredient:** deutetrabenazine

**Inactive ingredients:** ammonium hydroxide, black iron oxide, butyl alcohol, butylated hydroxyanisole, butylated hydroxytoluene, cellulose acetate, hydroxypropyl cellulose, hypromellose, isopropyl alcohol, magnesium stearate, polyethylene glycol, polyethylene glycol 3350, polyethylene oxide, polyvinyl alcohol, propylene glycol, shellac, sodium chloride, talc, titanium dioxide, FD&C blue #2 lake, and FD&C red #40 lake. The 6 mg and 12 mg extended-release tablets also contain FD&C yellow #6 lake.

**AUSTEDO:**

**Active ingredient:** deutetrabenazine

**Inactive ingredients:** ammonium hydroxide, black iron oxide, n-butyl alcohol, butylated hydroxyanisole, butylated hydroxytoluene, magnesium stearate, mannitol, microcrystalline cellulose, polyethylene glycol, polyethylene oxide, polysorbate 80, polyvinyl alcohol, povidone, propylene glycol, shellac, talc, titanium dioxide, and FD&C blue #2 lake. The 6 mg tablets also contain FD&C red #40 lake. The 12 mg tablets also contain FD&C yellow #6 lake.

Manufactured for:

**Teva Neuroscience, Inc.**

Parsippany, NJ 07054

©2023 Teva Neuroscience, Inc.

AUSMG-008

For more information, go to [www.AUSTEDO.com](http://www.AUSTEDO.com) or call 1-888-483-8279.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Revised: February 2023

AUS-45398